

{rokbox title=| Technical drawing of the bottom-line deployment system :: Image: Authors| thumb=|images/stories/ieo/imagenespublicaciones/centro-oceanografico-baleares-ieo-csic-trap-tools-monitoring-valls-2026-thumb-et-al.jpg|}images/stories/ieo/imagenespublicaciones/centro-oceanografico-baleares-ieo-csic-trap-tools-monitoring-valls-2026-et-al.jpg{/rokbox}

Maria Vigo, Ulla Fernandez-Arcaya, Joan Navarro, **Maria Valls**, **Beatriz Guijarro**, **M. Teresa Farriols**

Nixon Bahamón, Guiomar Rotllant, José A. García, Alba Rojas,

Enric Massutí

, Joan B. Company, 2026.

[Traps as tools for monitoring deep-sea ecosystems and promoting sustainable fishing](#)

. Marine Biology Research. Volume 22. 2026. Issue 3.

<https://doi.org/10.1080/17451000.2026.2621038>

Abstract: The Norway lobster (*Nephrops norvegicus*) and the deep-sea blue and red shrimp (*Aristeus antennatus*

) are among the most valuable fishing resources in the Mediterranean Sea. Both species are caught by bottom trawling, which causes significant ecological disturbance in deep-sea environments. This study investigates the viability of traps as a sustainable alternative for their capture. Experimental trap surveys were conducted in two geographical subareas (GSAs) of the western Mediterranean Sea: one targeting

N. norvegicus

in the GSA 06 at 350-550 m depth, and the other targeting

A. antennatus

in the GSA 05 at 550-850 m. We tested two bottom-line-trap designs and three trap types: for

N. norvegicus

, traps with 12 and 24 mm diamond mesh nets, and for

A. antennatus

, including also semi-floating traps with 24 mm diamond mesh. Traps were effective for

N. norvegicus

, particularly using 12 mm traps, although the finest mesh sizes increased bycatch of smaller individuals. In contrast,

A. antennatus

was not captured, potentially due to species-specific behavioural and operational constraints.

Other commercial species, such as

Plesionika spp.

, were effectively caught. Although traps are not a fully viable alternative to trawling, our findings highlight their potential as selective tools for research and monitoring in sensitive deep-sea habitats.

Keywords: Trap, deep-sea, *Nephrops norvegicus*, *Aristeus antennatus*, sustainable, fishery.