

{rokbox title=|Concentrations of plastic debris in surface waters of the global ocean :: Photo credit: Malaspina circumnavigation|  
thumb=|images/stories/ieo/imagenespublicaciones/centro-oceanografico-baleares-plasticos-malaspina-m-luz-fernandez-de-puelles-cozar-et-al-2014-thumb.jpg|}images/stories/ieo/imagenespublicaciones/centro-oceanografico-baleares-plasticos-malaspina-m-luz-fernandez-de-puelles-cozar-et-al-2014.jpg{/rokbox}

Andrés Cózar, Fidel Echevarría, J. Ignacio González-Gordillo, Xabier Irigoien, Bárbara Úbeda, Santiago Hernández-León, Álvaro T. Palma, Sandra Navarro, Juan García-de-Lomas, Andrea Ruiz, María L. Fernández-de-Puelles, and Carlos M. Duarte, 2014. [Plastic debris in the open ocean](#) . PNAS  
2014: 1314705111v1-201314705.

Abstract: There is a rising concern regarding the accumulation of floating plastic debris in the open ocean. However, the magnitude and the fate of this pollution are still open questions. Using data from the Malaspina 2010 circumnavigation, regional surveys, and previously published reports, we show a worldwide distribution of plastic on the surface of the open ocean, mostly accumulating in the convergence zones of each of the five subtropical gyres with comparable density. However, the global load of plastic on the open ocean surface was estimated to be on the order of tens of thousands of tons, far less than expected. Our observations of the size distribution of floating plastic debris point at important size-selective sinks removing millimeter-sized fragments of floating plastic on a large scale. This sink may involve a combination of fast nano-fragmentation of the microplastic into particles of microns or smaller, their transference to the ocean interior by food webs and ballasting processes, and processes yet to be discovered. Resolving the fate of the missing plastic debris is of fundamental importance to determine the nature and significance of the impacts of plastic pollution in the ocean.

Keywords: